

**Table 1.** Comparisons of whole carcass weight among regions by age class.

| REGION | AGE | WEIGHT | n    |
|--------|-----|--------|------|
| SO     | 1.5 | 100.3  | 1293 |
| EG     | 1.5 | 93.9   | 257  |
| NW     | 1.5 | 93.4   | 3098 |
| CE     | 1.5 | 90.7   | 4278 |
| NE     | 1.5 | 89.7   | 1484 |
| SO     | 2.5 | 116.5  | 544  |
| NW     | 2.5 | 113.6  | 2071 |
| CE     | 2.5 | 108.6  | 2789 |
| EG     | 2.5 | 108.4  | 170  |
| NE     | 2.5 | 104.6  | 984  |
| SO     | 3.5 | 132.8  | 237  |
| NW     | 3.5 | 127.7  | 753  |
| NE     | 3.5 | 117.0  | 543  |
| CE     | 3.5 | 116.5  | 1433 |
| EG     | 3.5 | 111.5  | 88   |

**Table 2.** Comparisons of antler beam length (in) among regions by age class.

| REGION | AGE | BEAM LENGTH | n    |
|--------|-----|-------------|------|
| SO     | 1.5 | 18.1        | 1745 |
| EG     | 1.5 | 17.9        | 1605 |
| CE     | 1.5 | 13.8        | 5354 |
| NE     | 1.5 | 13.7        | 2265 |
| NW     | 1.5 | 11.9        | 4254 |
| SO     | 2.5 | 28.0        | 797  |
| NW     | 2.5 | 26.7        | 3323 |
| CE     | 2.5 | 25.2        | 3561 |
| EG     | 2.5 | 24.6        | 1205 |
| NE     | 2.5 | 23.2        | 1569 |
| SO     | 3.5 | 35.2        | 327  |
| NW     | 3.5 | 33.9        | 1179 |
| CE     | 3.5 | 30.4        | 1796 |
| NE     | 3.5 | 30.0        | 878  |
| EG     | 3.5 | 28.3        | 553  |